



## **Polyploidy can confer superiority to West African *Acacia senegal* (L.) Willd. trees**

Diallo, Adja Madjiguene; Nielsen, Lene Rostgaard; Kjær, Erik Dahl; Petersen, Karen K.; Ræbild, Anders

*Published in:*  
Frontiers in Plant Science

*DOI:*  
[10.3389/fpls.2016.00821](https://doi.org/10.3389/fpls.2016.00821)

*Publication date:*  
2016

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Diallo, A. M., Nielsen, L. R., Kjær, E. D., Petersen, K. K., & Ræbild, A. (2016). Polyploidy can confer superiority to West African *Acacia senegal* (L.) Willd. trees. *Frontiers in Plant Science*, 7, 821.  
<https://doi.org/10.3389/fpls.2016.00821>

## Polyploidy can confer superiority to West African *Acacia senegal* (L.) Willd. trees

Adja M. Diallo<sup>1,2</sup>, Lene R. Nielsen<sup>1</sup>, Erik D. Kjær<sup>1</sup>, Karen K. Petersen<sup>3</sup>, Anders Ræbild<sup>1\*</sup>

<sup>1</sup>Department of Geosciences and Natural Resources, University of Copenhagen, Denmark, <sup>2</sup>Centre National de Recherches Forestières, Institut Sénégalais de Recherches Agricoles, Senegal, <sup>3</sup>Department of Food Science, University of Aarhus, Denmark

*Submitted to Journal:*  
Frontiers in Plant Science

*Specialty Section:*  
Functional Plant Ecology

*ISSN:*  
1664-462X

*Article type:*  
Original Research Article

*Received on:*  
08 Mar 2016

*Accepted on:*  
26 May 2016

*Provisional PDF published on:*  
26 May 2016

*Frontiers website link:*  
[www.frontiersin.org](http://www.frontiersin.org)

*Citation:*  
Diallo AM, Nielsen LR, Kjær ED, Petersen KK and Ræbild A(2016) Polyploidy can confer superiority to West African *Acacia senegal* (L.) Willd. trees. *Front. Plant Sci.* 7:821. doi:10.3389/fpls.2016.00821

*Copyright statement:*  
© 2016 Diallo, Nielsen, Kjær, Petersen and Ræbild. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Provisional

**Polyploidy can confer superiority to West African *Acacia senegal* (L.)  
Willd. trees**

Adja Madjiguene Diallo<sup>1,2\*</sup>, Lene Rostgaard Nielsen<sup>1</sup>, Erik Dahl Kjær<sup>1</sup>, Karen Koefoed Petersen<sup>3</sup> and  
Anders Ræbild<sup>1</sup>

<sup>1</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen,  
Denmark

<sup>2</sup>Centre National de Recherches Forestières/Institut Sénégalais de Recherches Agricoles  
(CNR/ISRA), Dakar, Senegal

<sup>3</sup>Department of Food Science, Aarhus University, Aarslev, Denmark

**\*Correspondence:**

Anders Ræbild

are@ign.ku.dk

**Running head:** Superiority of polyploid *Acacia senegal*

**Number of figures and tables:** 7

**Number of words:** 7953

21 **Abstract**

22 Polyploidy is a common phenomenon in the evolution of angiosperms. It has been suggested that  
23 polyploids manage harsh environments better than their diploid relatives but empirical data  
24 supporting this hypothesis are scarce, especially for trees. Using microsatellite markers and flow  
25 cytometry, we examine the frequency of polyploids and diploids in a progeny trial testing four  
26 different populations of *Acacia senegal*, a species native to sub-Saharan regions of Africa. We  
27 compare growth between cytotypes and test whether polyploid seedlings grow better than diploids.  
28 Our results show that polyploids coexist with diploids in highly variable proportions among  
29 populations in Senegal. *Acacia senegal* genotypes were predominantly diploid and tetraploid, but  
30 triploid, pentaploid, hexaploid and octaploid forms were also found. We find that polyploids show  
31 faster growth than diploids under our test conditions: in an 18 years old field trial, polyploid  
32 superiority was estimated to be 17% in trunk diameter and 9% in height while in a growth chamber  
33 experiment, polyploids grew 28 % taller, but only after being exposed to drought stress. The results  
34 suggest that polyploid *A. senegal* can have an adaptive advantage in some regions of Africa.

35 **Key words:** adaptation - arid zone trees - drought stress – flow cytometry - microsatellite markers -  
36 morphological differentiation - *Senegalia senegal*

Provisional

## 1. Introduction

Polyploidy, the achievement of more than two sets of chromosomes through gametic non-reduction and to a lesser degree somatic doubling has important ecological and evolutionary consequences for speciation (Madlung, 2013). In nature, polyploidy arises via intraspecific genome doubling (autopolyploidy) or merging of genomes of distinct species through hybridization and chromosome doubling (allopolyploidy) (Stebbins, 1950). Most angiosperms are believed to have undergone one or more polyploidization events (Soltis *et al.*, 1999). It has been estimated that polyploids form at the frequency of approximately 1 per 100,000 individuals (Ramsey & Schemske, 1998; Levin, 2002) and that 2 – 4 % of all speciation events involve polyploidization (Otto & Whitton, 2000). The high level of polyploidization in the evolutionary history of flowering plants suggests that polyploidy plays an important role in adaptive evolution of plants in natural populations (Van de Peer *et al.*, 2009). Successful polyploidization is generally accompanied by morphological, phenological, physiological, and ecological changes in plants (Levin, 2002), and may produce individuals that can tolerate fluctuating environments (Soltis *et al.*, 2004; Prentis *et al.*, 2008), make use of new niches or by other means become more successful than their progenitor species (Leitch & Leitch, 2008).

In theory, local co-occurrence of intraspecific cytotypes is evolutionary unstable, driving the minority cytotype towards extinction unless cytotypes have different ecological preferences or strong prezygotic barriers are present between ploidy levels (Husband 2000, Kennedy *et al.*, 2006). Nevertheless, co-distribution of individuals belonging to different ploidy levels in heteroploid species (species with different levels of ploidy) is not uncommon and has been reported in *e.g.* *Solidago altissima*, *Ranunculus parnassifolius* and *Centaurea phrygia*, *C. stoebe* (Halverson *et al.*, 2008; Cires *et al.*, 2010; Koutecky *et al.*, 2012). With the recent introduction of flow cytometry it is now possible to explore the cytotype dynamics in species with mixed ploidy levels in terms of cytotype distribution, hybridization and segregation.

Morphological changes in polyploids include a general increase of cell size with increased levels of ploidy (*e.g.* Kudo & Kimura 2002), sometimes leading to changes in the dimensions of plants, such as larger leaf, flower and fruit sizes compared to diploids (Maherali *et al.* 2009, Pettigrew *et al.* 2012). Also micro-morphological changes occur, including larger but more dispersed stomata in polyploids than in diploids (Mishra 1997, Pettigrew *et al.* 2012). Such changes are likely to affect plant environment interactions, for example through modification of gas exchange. It was earlier suggested that polyploids withstand harsh environments like subarctic regions, high elevations and xeric environments better than diploids (Love & Love, 1949) perhaps due to their higher levels of heterozygosity and genetic diversity (Lowry & Lester, 2006). This was supported for example by observations of higher colonization potentials in polyploids, increased frequencies of polyploids from warmer to colder latitudes (Manton, 1934, Hagerup, 1939, Brochmann *et al.*, 2004, Lowry & Lester, 2006) and larger ecological amplitude of polyploids compared to diploids (*e.g.* Liu *et al.*, 2011, Schlaepfer *et al.*, 2010).

Recent studies have found that polyploid cytotypes in heteroploid species complexes of herbaceous plants have better drought tolerance than their diploid progenitors (*Chamerion angustifolium*, Maherali *et al.*, 2009; *Brachypodium distachyon*, Manzaneda *et al.*, 2012), although Buggs & Panell (2007) found that diploid *Mercurialis annua* performed better than hexaploids across a range of natural sites and that the cytotypes did not differ in performance under drought stress. Increased drought tolerance may be related to higher resistance towards cavitation in the xylem as discovered in *Atriplex canescens*, a shrub species from the deserts of Southwestern U.S. (Hao *et al.*, 2013). Nevertheless, it is still discussed under which circumstances polyploidy confer higher fitness (Madlung, 2013), and a comparison of many North American diploid and polyploid species showed no significant differences in extent of range or geographical distribution between ploidy levels

(Martin & Husband, 2009). Unfortunately very few studies showed the relative performance of diploids and polyploids under controlled conditions (Soltis *et al.* 2010).

Variation in ploidy level within species is also known from trees (e.g. the *Adansonia digitata*/*A. kilima* complex, Pettigrew *et al.*, 2012, *Betula papyrifera*, Li *et al.*, 1996, *Populus tremula*, Johnsson 1940, *Acacia mearnsii*, Beck *et al.* 2003). Because increased cell size is likely to influence hydraulic properties via the influence on conduits (see e.g. Hao *et al.*, 2013, Maherali *et al.*, 2009), studies on trees with their massive xylem, large size and long potential exposure to climatic extremes are particularly interesting. Yet there are few studies comparing performance of trees with different ploidy levels, and the studies that exist focus on short-term responses of seedlings to stress (Li *et al.*, 1996, Li *et al.*, 2009). Studies of performance of mature polyploid versus diploid trees are to our knowledge absent.

Recently it was discovered that *Acacia senegal* (L.) Willd. exists in different levels of ploidy. The species grows naturally in the semi-arid sub-Saharan regions of Africa as well as in India and Pakistan and plays an important role in agroforestry systems by providing fuel, fodder for livestock and restoring soil fertility besides producing Gum Arabic. Gum Arabic is a natural exudate collected from branches and stems after tapping during the dry season, and is only produced when the species is grown under dry conditions (Wekesa *et al.*, 2009). The gum provides an important income for rural people. Most of the populations across the distribution area seem to be composed of diploid trees, but tetraploid individuals ( $2n=4x=52$ ) were discovered in populations from Mali, Sudan and Ethiopia (Assoumane *et al.*, 2013). Due to a limited sample size within each population the ratio between diploid and polyploid individuals was not resolved. Based on chloroplast data, the authors suggest that polyploid *A. senegal* is allopolyploid (Assoumane *et al.*, 2013). The parent species that hybridized with diploid *A. senegal* may have been *Acacia laeta* reported to be a triploid hybrid ( $3x = 39$ ) between *A. senegal* and *A. mellifera* (Ross, 1979), but the origin and type of polyploidy in *A. senegal* is still not verified.

Investigating a progeny trial with mature *A. senegal*, we discovered that the trees had mixed ploidy levels (Diallo *et al.*, 2015). As the trial was established with the purpose of breeding for increased gum production, the trees were planted in an experimental design and thus represent a unique possibility for studying the long-term performance of di- and polyploid trees. In this paper, we specifically 1) explore the distribution of cytotypes in trees originating from four different locations, and in their corresponding offspring, 2) compare the growth performance of trees with different ploidy levels, 3) compare the growth of seedlings with different ploidy level under drought stress and 4) compare the morphology of plants with different ploidy levels.

## 2. Materials and methods

### 2.1. Study species

*Acacia senegal* (L.) Willd. is a multipurpose tree that belongs to the family Fabaceae. Recently, it has been suggested to transfer the species to the new genus *Senegalia* (Maslin, 2006). Here we maintain the rule of first priority and thus the name *Acacia senegal*. The species is described as consisting of the four varieties *senegal*, *kerensis*, *rostrata* and *leiorhachis* based on differences in inflorescence axis, pod and tree shape and phenology (Odee *et al.*, 2012). Only *A. senegal* var. *senegal* has been reported in Senegal.

### 2.2. The field trial

137 The plant material of *A. senegal* used in this study originated from a progeny trial in Senegal  
138 established in 1994 in Dahra, Senegal (15° 20' N and 15° 28' W). The annual precipitation at the site  
139 is approximately 410 mm, and the annual mean temperature is 27°C (climate data estimated from  
140 Worldclim based on the 1950-2000 period, see Hijmans *et al.* 2005). The soil at the site is sandy, and  
141 the natural vegetation in the area consists of mainly *Acacia tortilis* and *Balanites aegyptiaca*  
142 (Pontanier *et al.* 2003).  
143

144 In November and December 1993, seeds were collected from four populations (provenances)  
145 representing the natural distribution area of the species in Senegal (Fig. 1): Ngane, located in the  
146 centre of Senegal and characterized by saline soils and 620 mm of rainfall; Diamenar, located in a  
147 dryer region in the north with 300 mm of rainfall; Daiba and Kidira, located in the north-eastern and  
148 south-eastern parts with 430 mm and 600 mm of rainfall, respectively. At each site seeds were  
149 collected from 15 trees considered to have desirable phenotypes based on superior health, trunk  
150 diameter, crown diameter and height. In the natural stands diploids and polyploids are  
151 indistinguishable with the naked eye, implying no bias for or against any type during seed  
152 collections. Seeds were kept in separate lots for each mother tree, thus giving 60 seedlots, and in  
153 1994 seeds were pretreated in sulfuric acid (98 %) for 6 min and sown in polyethylene bags with  
154 nursery soil. Two seeds were sown per bag, and in cases where both seeds germinated, one of the  
155 seedlings was randomly selected and removed. Seedlings were raised in a nursery and in August  
156 1994 (during the rainy season), 30 healthy seedlings per seedlot were selected and planted at the  
157 Dahra site. Prior to plantation, weeding and clearing were undertaken in the site. The trial was  
158 established in a randomized complete single tree block design with all seedlots represented by one  
159 tree in each block, replicated thirty times (30 blocks). The initial number of plants was thus  $4 \times 15 \times$   
160  $30 = 1800$  trees. Trees were spaced 5 x 5 meters from each other.  
161

162 One year after planting, in August 1995, survival and height of all trees were assessed. Height was  
163 measured as vertical height from the ground to the top of the tree. In February 2012, a second  
164 assessment was conducted on the 634 surviving trees. The maximum vertical height was determined  
165 using a height rod, trunk diameter at 30 cm from the soil surface was assessed using a diameter tape,  
166 and crown diameter was estimated as the average of two perpendicular measures of the edge of the  
167 crown projected to the ground. In 2012, cambium samples were taken from every tree for assessment  
168 of the ploidy level (see later).  
169

### 170 2.3 Growth chamber experiment

171 In November 2012, 108 pods were collected from 76 parent trees in the field trial. The parent trees  
172 were selected to cover all four sites of origin and – to the extent possible – different levels of ploidy  
173 in all sites of origin. One pod was randomly chosen from each tree except for trees from Ngane  
174 where 1-6 pods were collected per tree (due to higher frequency of polyploidy in this provenance as  
175 described below).  
176

177 The number of seeds per pod was counted and seed dimensions (length and width) were measured  
178 using a Vernier caliper. Seeds were pretreated with sulfuric acid (95-97 %) for 10 min to release seed  
179 dormancy, kept under sterile conditions in a laminar hood (Thermo Scientific, SAFE 2020, Germany)  
180 and germinated in boxes containing sterilized vermiculite and incubated at 29 °C and 16 h  
181 photoperiod in a growth chamber. The germination rate was registered, and the ploidy level of all  
182 seedlings was determined (see below).  
183

184 To compare the growth between different levels of ploidy, 132 seedlings were transplanted in peat  
185 soil (Plugg och Såjord) from Weibulls Horto AB, Sweden. 500 g of soil was filled in 17 plastic boxes  
186 each. The seedlings consisted of 83 diploids, 46 tetraploids and three hexaploids. Each box was



187 considered as one block and contained eight seedlings representing different sites of origin  
188 (provenance), descendance and ploidy levels. By descendance we here understand all seedlings  
189 descending from a single mother tree in the Dahra field trial. Because of the restricted number of  
190 polyploid seedlings in some sites of origin (provenances), the design was imbalanced, but each box  
191 contained at least three sites of origin and one set of diploids and polyploids from the same site of  
192 origin. For example, block no. 1 included four tetraploids and one diploid from the Ngane origin, and  
193 one diploid of each of the Daiba, Diamenar and Kidira origins, while block no. 17 contained one  
194 diploid and one tetraploid of the Kidira origin, and three diploids of each of the Daiba and Diamenar  
195 origins. Despite the imbalance in the trial, this design allowed us to assess the growth of the different  
196 levels of ploidy because there was always at least one pair of diploid and polyploid seedlings from the  
197 same origin in each block. We planted seedlings in boxes to make sure that all plants in a box were  
198 exposed to the same water level irrespective of plant size, leaf area and stomatal conductance  
199 (Verslues et al. 2006).

200  
201 The initial weight of seedlings was recorded before transplanting into soil. We first compared the  
202 growth under well-watered conditions (85% of field capacity) for five weeks in the growth chamber  
203 at 16 h photoperiod. The temperature varied between 28 and 33 °C, while relative air humidity ranged  
204 between 47 and 71%. Boxes were weighed every day and the amount of water lost by  
205 evapotranspiration was added to maintain 85% of field capacity. Plant height was measured weekly  
206 from the soil surface to the apical bud, and the numbers of leaves and branches were counted after  
207 three weeks. Leaf length, leaflet length and width were measured after five weeks of growth under  
208 well-watered conditions. Likewise, stomatal size and density were determined on 50 plants (diploid  
209 and polyploid) and two randomly chosen leaflets per seedling. Leaflets were stained with Toluidine  
210 Blue O dissolved in 0.05 % of benzoate buffer and water at pH 4.4 (O'Brien & McCully, 1981) and  
211 viewed with a microscope (Olympus Cx40 RF 200, Japan) under x 40 magnification. Two  
212 microscopic grids were examined per leaflet, totaling 200 counts. To determine the size of stomata,  
213 20 random individual stomata per seedling were measured (scale bar 100 µm) at magnification x 40  
214 and the mean stomatal length and density was calculated.

215  
216 After five weeks, drought stress was applied by reducing the amount of added water to 47% of field  
217 capacity. In a pilot study, this was shown to be close to the wilting point. Again, in order to maintain  
218 the field capacity at 47%, boxes were weighed every day and the amount of water lost was added.  
219 Plant height was recorded weekly for six weeks and the fresh and dry biomasses of seedlings (after  
220 drying at 80°C for 48 h) were assessed at the end of the experiment.

221

222

## 223 2.4. Ploidy level assessment

224

225 For 59 trees from the Dahra field trial, twigs of 20 cm length with vegetative buds were collected in  
226 April 2014 and placed in a lab with their proximal end in water. After two weeks of forcing, one  
227 complete leaf was collected from each twig and analyzed by flow cytometry. Seeds descending from  
228 the mother trees in the Dahra field trial were germinated as described above, and flow cytometry was  
229 performed on 3 weeks old leaves in the lab.

230

231 Flow cytometry was performed using a Partec PA II flow cytometer equipped with an HBO-100  
232 mercury arc lamp (Partec GmbH, Germany) and filter combination for DAPI staining (Partec 06-03-  
233 310). Fresh leaf samples of two-weeks-old plants or from forced twigs were chopped for 30 s in a  
234 petri dish containing 0.6 ml of Citric acid buffer and left for 5 min to allow nuclei release. The nuclei  
235 were stained by adding 2.5 ml of fluorescent solution containing 5 µM DAPI (4,6-Diamino-2-  
236 phenylindol dihydrochlorid) and left for another 5 min (Otto, 1990). The suspension of nuclei was

passed through a nylon filter with pore size 50 µm to remove large debris. The DAPI binds to the A-T bases of DNA and the intensity of the fluorescence emitted will reflect the number of bounds and therefore also the DNA content in such DAPI-labelled nuclei. The relative fluorescence of total DNA of single nuclei was analyzed and in each sample the DNA content of 5000 nuclei was checked. Samples of *Miscanthus sinensis* with known ploidy (diploid) was used as an internal standard. The standard produced two peaks: a major peak corresponding to 2x DNA quantities of the majority of the cells and a minor peak corresponding to 4x DNA quantities from cells in mitotic interphase.

The gain was adjusted so that the peak of diploid *A. senegal* was localized on channel 50 corresponding to one large (2C) and one small (4C) peak. Plants were regarded as tetraploids if histograms showed one major 4C peak, a small 8C peak and no 2C peak.

To assess the exact DNA content of the genome, a subset of 26 leaf samples from 13 DAPI-examined plants (3 diploids, 3 triploids, 3 tetraploids, 1 pentaploid and 3 hexaploids) was analyzed using flow cytometry with propidium iodide dye according to the protocol by Doležal *et al.* (2007).

All 634 living trees in the field trial were genotyped with 8 polymorphic microsatellite markers (SSR) as described in Diallo *et al.* (2015). By comparing the flow cytometry results with the SSR markers for the 59 mature trees, we concluded that polyploids could always be separated from diploids by the presence of more than two alleles per locus in at least 1 of the 8 loci (Appendix 1). Based on their SSR genotypes, we therefore assigned all 634 trees in the field trial to either diploid or polyploid status, but not distinguishing between tri-, tetra-, penta- or hexaploids.

## 2.5. Statistics

For the data from the field trial, a generalized linear analysis of variance was applied to test differences between the growth of diploid and polyploid trees in the trial. Trees that were assessed in 1995 but were not alive in 2012 were excluded, as their ploidy level was unknown. The analysis included the effects of ploidy level (diploid or polyploid), site of origin (Daiba, Diamenar, Kidira or Ngane) and block (30 levels) according to the following model:

$$Y = \text{Ploidy} + \text{Site of origin} + \text{Block} + \text{Error} \quad (1)$$

Where the effects of ploidy and site of origin were considered as fixed, block was considered as random, and the error followed a normal distribution with expectation zero. The average performance of diploids and polyploids was estimated as least square means from the analysis, *i.e.* averages corrected for systematic differences among provenances.

Data from the growth chamber experiment were analyzed in two steps. The first step was a model with the effects of descendance and blocks:

$$Y = \text{Descendance} + \text{Block} + \text{Error} \quad (2)$$

Where descendance was considered a fixed effect and the block was considered as random. From this model, we calculated the least square mean values of all variables for each descendance. Due to the limited number of samples for stomatal density and length, calculation of least square means was not possible and instead the mean values were calculated. Mean values were also calculated for seed traits. Next, as each descendance was either diploid or polyploid and originated from one of the four sites of origin, we applied the following model:

$$Y = \text{Ploidy} + \text{Site of origin} + \text{Error} \quad (3)$$

Where Y denotes the least square means (or means) of the descendant families from model (2), and ploidy and site of origin were considered as fixed effects.

All analyses were performed using the GLM procedure in the SAS 9.3 Software (SAS Institute Inc. 2014). Assumptions of variance homogeneity and normality of residuals were accepted for all the studied characters based on visual inspection of residual plots. However, for the ratio between fresh and dry weight in the growth chamber experiment a single outlier (Kidira family 7) was identified and deleted from the data.

### 3. Results

#### 3.1. Ploidy level

Both diploid and polyploid individuals were found among trees from all four origins, but at very different frequencies: 136 of 164 trees (83%) originating from Ngane were polyploid, compared to 3 of 178 trees in Diamenar (2%), 14 of 146 trees in Daiba (10 %) and 16 of 117 trees in Kidira (14%).

Flow cytometric analyses of seedlings allowed separation between diploid, triploid, tetraploid, pentaploid and hexaploid individuals corresponding to a mean 2C DNA content of  $1.25 \pm 0.02$  pg,  $1.96 \pm 0.05$  pg,  $2.60 \pm 0.03$  pg, 3.19 pg and  $3.83 \pm 0.08$  pg respectively. All offspring from diploid mothers were diploid, while tetraploid mothers produced either offspring with the same ploidy level (tetraploid) or higher levels (pentaploid, hexaploid and octaploid). Seeds from the same pod collected on the tetraploid mother NG16\_B19 gave rise to one tetraploid and one hexaploid seedling, while all seedlings coming from the tetraploid mother DA15\_B3 were hexaploid. Of 13 offspring tested from the tetraploid parent NG10\_B8, 11 were tetraploid, one was pentaploid and one was hexaploid; of the 16 plants examined from the tetraploid mother NG20\_B3, one was octaploid and 15 were tetraploid. Out of three seedlings tested from the triploid individual DA1\_17, two were triploid and one was tetraploid (Appendix 1).

#### 3.2. Growth differences between diploid and polyploid trees in the progeny trial

The polyploid trees in the Dahra trial were significantly taller than the diploid trees 1 year after planting in 1995. In 2012, after 18 years, differences between diploid and polyploid trees for both height and trunk diameter were still significant, whereas crown diameter did not differ between ploidy levels. The superiority of polyploids compared to diploids based on least square mean estimates after 1 year was 18% for height, while 9% for height and 17% for trunk diameter after 18 years. The true differences between average of diploid and polyploid trees (i.e not corrected for systematic effects of provenances) were substantially larger: 49% for height at age 1; and 15% and 24% for height and diameter respectively at age 18 (Table 1, Fig. 2).

#### 3.3. Phenotypic differences between diploid and polyploid seedlings in the drought stress test

Polyploids differed significantly from diploids in seed length and width, initial fresh weight (at week 0 when transplanted to the boxes), leaflet length, stomatal density and length. Heights were similar until weeks 10 and 11. Total fresh weight was borderline significant at the end of the trial, and the fresh weight / dry weight ratio differed significantly between ploidy levels (Table 2, Fig. 2).

The morphological parameters showed that the polyploids tended to be larger than the diploids (Table 2). Seed length and width were 12% and 10% larger in polyploids compared to diploids, respectively. Polyploids had leaflets that on average were 19% longer than in diploid individuals, and the differences in stomatal density and length were pronounced: polyploid individuals were characterized by 54% wider stomata but with lower stomatal density (31% less) compared to diploids.

Prior to drought stress, the plant height was similar for both ploidy levels, but after 10 and 11 weeks (corresponding to 5 and 6 weeks of water deficit), the tetraploids has grown taller than diploids (22 and 28 %, respectively) (Table 2, Fig. 3). At the end of the trial, polyploid *A. senegal* seedlings had 60% larger fresh weight than diploids. Differences in dry weight were smaller because the fresh weight / dry weight ratio was 9% larger in polyploids than in diploids.

## 4. Discussion

### 4.1. Evolution of polyploidy in *A. senegal*

Our results showed that *A. senegal* can occur in more than two levels of ploidy, which supplement the results of Assoumane *et al.* (2013) and Odee *et al.* (2015) who reported diploid and tetraploid individuals. In our study, we found that diploids were most frequent followed by tetraploids. Pentaploids, triploids, and hexaploids were also present among seedlings although in small quantities. Based on small sample sizes, Odee *et al.* (2015) showed co-existence of ploidy types. This result is qualified by the present study, where we show that the frequency of polyploid individuals can vary significantly among natural populations in Senegal. Polyploids were dominant in the population from the central Senegal (Ngane) characterized by saline soils, whereas higher proportions of diploids were found in Diamenar (North), Daiba (North-east) and Kidira (South-east). Nevertheless, a few diploid individuals from the saline site (Ngane) were also identified and polyploids occurred in low frequency in non-saline areas (Kidira, Diamenar and Daiba sites).

Our data comparing the ploidy level of trees and their offspring indicated only very limited hybridization between cytotypes, even in a setup where diploid and polyploid trees were grown side by side in a field trial. Further, offspring from diploid mothers were always diploid suggesting that hybridization between cytotypes with diploid maternal trees must be very rare if at all possible. The higher ploidy levels occasionally found in offspring from tetraploid mothers on the other hand suggest that tetraploid mothers rather frequently produce some gametes that are unreduced. Hexaploid seedlings from tetraploid mothers may have been formed by unreduced egg cells (4n) sired by reduced pollen gametes from a tetraploid pollen donor. The identified pentaploid seedling could have been formed by a pollination event involving a hexaploid pollen donor from the field trial (however not among the trees that were tested with FCM). Alternatively the pentaploid seedling could originate from a fusion between an unreduced egg cell from the tetraploid mother and a reduced pollen gamete from a diploid pollen donor i.e. reflecting cytotype hybridization. In relation to this aspect, we found a single triploid mother DA1\_B17 which might have been formed by cytotype hybridization in the previous generation. This triploid tree produced both triploid and tetraploid seedlings. Triploids have previously been reported when conspecific diploids and tetraploids co-occur in the same area as *e.g. Chameron angustifolium* (Husband, 2004) and may play an important role as a triploid-bridge allowing gene flow through mating between diploid and polyploid individuals (Henry *et al.*, 2005) with recurrent polyploid formation in the population.

The frequency of triploid seedlings (2 out of 162 – both from a triploid mother) observed in our study is, however, low compared to the reported frequencies of triploids in species with mixed populations

(range of 2 - 22 %) as reviewed by Soltis *et al.* (2010). Also, no triploid seedlings were observed from either diploid or tetraploid mother trees, which support the presence of a significant reproductive barrier. Stebbins (1971) predicted that mating between ploidy levels is likely only from diploid fathers to polyploid mothers. Unidirectional mating from diploids to tetraploids is known from other species complexes such as *Sorghum* (diploid *Sorghum bicolor* and tetraploid *Sorghum halepense*) (Arriola & Ellstrand 1996), *Capsella rubella* (to the allotetraploid descendant *C. bursa-pastoris*) (Slotte *et al.*, 2008), and *Arabidopsis arenosa* (Jørgensen *et al.*, 2011). As pollen is aggregated in polyades in *A. senegal* and the stigmatic cavity is cup-shaped (Tandon *et al.*, 2001) morphological size differences between cytotypes could also restrict hybridization. Additional detailed studies are needed to clarify the strength of the reproductive barriers between cytotypes of *A. senegal*, and if pollination is always unidirectional under natural conditions.

#### 4.2. Adaptive potential and evolutionary success of polyploids

Polyploid trees often occupy drier habitats than their diploid relatives (e.g. Li *et al.*, 1996, Pettigrew *et al.*, 2012). Experiments have shown different performance of diploids and polyploids under drought stress, but are limited by their short duration and experimental setup (Li *et al.*, 1996, Li *et al.*, 2009). Unfortunately, there is very limited evidence from long-term field trials on the relative performance of trees with different ploidy levels. The faster growth of polyploid *A. senegal* in our study is to our knowledge the first observation of superiority of mature natural polyploid trees in a field trial and indicates that at least under some conditions, trees with high ploidy levels will have an adaptive advantage. Although the effects of ploidy level and origin were to some extent confounded due to the observed unequal distribution of polyploids, our statistical analysis showed that the positive effect of being polyploid remained even when accounting for the effect of origin.

The field trial represents relatively dry conditions close to the Northern limit of distribution of the species towards the Sahara desert. Since the relative performance of diploid and polyploid *A. senegal* has not been investigated under wetter conditions, it is not possible to conclude whether the better growth is found only under dry conditions or indicates a general superior performance.

Still, the growth chamber experiment showed that polyploid plants only grew faster than diploids after the plants were subjected to water stress. The limited number of polyploid plants did not allow us to include a control treatment where seedlings were continuously raised without water stress, and we hence do not separate potential ontogenetic effects from effects of drought. The question of whether the adaptive advantage of polyploids is limited to dry conditions or applies over a broader range of environments therefore remains unresolved. Reciprocal experiments ('optimal' versus a single abiotic stress factor) to test the relative performance of di- and polyploids under different stress situations are needed to conclude whether polyploids are generally superior or if it is only the case under dry conditions (*cf.* Soltis *et al.*, 2010). Investigations in other heteroploid tree species or species complexes are needed to reveal if the observed effects of polyploid in *A. senegal* reflect a so-far undiscovered pattern in trees species growing under stressful conditions in Sahel.

Polyploidy is often associated with a difference in plant phenotype (e.g. increase in cell size, enlarged floral structure, pollen, stomata and robust stems) when compared to the diploid relatives (Ramsey & Schemske 1998; Madlung, 2013). In our study, we found the first evidence of phenotype differentiation between cytotypes in *A. senegal*, as seeds, leaflets and stomata were larger and stomatal densities were smaller in tetraploids of *A. senegal*. This confirms results reported for other tree species, such as *Adansonia digitata* (Pettigrew *et al.*, 2012), *Betula papyrifera* (Li *et al.*, 1996), *Acacia mangium* (Harbard *et al.*, 2012) and *Acacia maerensii* (Beck *et al.*, 2003). Studies based on neopolyploids have shown that many of these polyploid characteristics are directly linked to

increased genome size (e.g., Harbard *et al.*, 2012). It is unknown whether the size differences in *A. senegal* are caused by increased genome size, increased genetic diversity or a combination of both.

The observed phenotypic differences are likely to lead to differences in physiology, as gas movement in and out of leaves is affected by leaf size, size and distribution of stomata. It has been hypothesized that the fewer, but larger stomata observed in polyploids can change stomatal conductance and confer increased water use efficiency to polyploids under drought stress (e.g. Li *et al.*, 1996; Li *et al.*, 2009; Pettigrew *et al.*, 2012). Assuming that width and depth of the stomatal pore is proportional to the length, it can easily be estimated following Franks & Farquahar (2001) and Franks & Beerling (2009)) that stomatal conductance is expected to be 5% larger in polyploids than diploids. On the other hand, estimates based on leaf dimensions (Nobel, 2009) suggest that leaf boundary layer conductance will be reduced by 9% in polyploids compared to diploids. Hence the effects of leaf size tend to negate effects of changed stomatal size and density, and the expected overall effects of ploidy level on water use efficiency are therefore unclear. Detailed anatomical studies, coupled with assessments of gas exchange on trees with different levels of ploidy will be important in order to infer on potential mechanisms behind the putative selective advantage of polyploidy in *Acacia senegal*. For example, in the herbaceous perennial *Chamerion angustifolium*, Maherali *et al.* (2009) found that tetraploids characterized by large stomata and wide xylem vessels did not differ from diploids in stomatal conductance and gas exchange when grown under drought conditions. However, increased hydraulic conductivity was believed to cause increased drought resistance of tetraploids in this species.

Another observed phenotypic difference with potential physiological consequences is the larger fresh to dry weight ratios of polyploids compared to diploids (Table 2) causing the ploidy levels to differ almost significantly in fresh weight, but not in dry weight. High water contents indicate either a larger capacity for osmotic adjustment or an increased elasticity of cell walls (Verslues *et al.* 2005), and Li *et al.* (1996) suggested that polyploids might have a larger ability to adjust their osmotic potential under drought stress. If this is indeed the case, it may explain part of the better performance of polyploids in *A. senegal*.

In conclusion, the co-existence of different ploidy levels in natural populations was confirmed while the pattern of segregation supports that gene flow between cytotypes is limited. Our results document increased growth of polyploid *A. senegal* both in the field trial and under growth chamber conditions, but it remains to be verified if superiority of polyploids is expressed only under relatively dry conditions or applies more generally.

#### Author contributions

AD, LN, EK and AR conceived the ideas; AD, LN and KP collected the data; AD, EK and AR carried out the statistical analyses; all authors analyzed and interpreted the data, and all authors contributed to writing of the paper.

#### Funding

This research is funded by the Islamic Development Bank (IDB) under the PhD Merit Scholarship program (Student Grant Number 600014414), by ISRA, Senegal, and by the University of Copenhagen, Denmark.

#### Conflict of interest

483 The authors declare no conflict of interest.

484

#### 485 **Acknowledgements**

486

487 The authors wish to thank Momar Wade for assistance with fieldwork, Lene Hasmark Andersen for  
488 genotyping support, Karin Henriksen for assisting with the flow cytometric analysis, Hanne N.

489 Rasmussen for her support in the study of stomata and Antoine Sambou for data gathering. We are  
490 grateful to Richard Buggs for his comments on a previous version of the manuscript, and thank the  
491 anonymous reviewers for their constructive comments.

Provisional

## References

- Arriola, P. E., Ellstrand, N. C. (1996). Crop-to-weed gene flow in the genus *Sorghum* (Poaceae): Spontaneous interspecific hybridization between Jonsongrass, *Sorghum halepense*, and crop sorghum, *S. bicolor*. *Curr. Opin. Plant Biol.* 8, 135-141.
- Assoumane, A., Zoubeirou. A. M., Rodier-Goud, M., Favreau, B., Bezançon. G., Verhaegen, D. (2013). Highlighting the occurrence of tetraploidy in *Acacia senegal* (L.) Willd. and genetic variation patterns in its natural range revealed by DNA microsatellite markers. *Tree Genet. Genomes* 9, 93-106.
- Beck, S. L., Dunlop, W. R., Fossey, A. (2003). Stomatal length and frequency as a measure of ploidy level in black wattle, *Acacia mearnsii* (de Wild). *Bot. J. Linn. Soc.* 141, 177-181.
- Buggs, R. J., Pannell, J. R. (2007). Ecological differentiation and diploid superiority across a moving ploidy contact zone. *Evolution* 61, 125-140.
- Brochmann, C., Brysting, A. K., Alsos, I. G., Borgen, L., Grundt, H. H., Scheen, A. C., Elven, R. (2004). Polyploidy in arctic plants. *Biol. J. Linn. Soc.* 82, 521-536.
- Cires, E., Candela, C., Revilla, M. Á., Fernández Prieto, J. A. (2010). Intraspecific genome size variation and morphological differentiation of *Ranunculus parnassifolius* (Ranunculaceae), an Alpine–Pyrenean–Cantabrian polyploid group. *Biol. J. Linn. Soc* 101, 251-271.
- Diallo, A.M., Nielsen, L.R., Hansen, J.K., Ræbild, A., Kjær, E.D. (2015): Study of quantitative genetics of gum arabic production complicated by variability in ploidy level of *Acacia senegal* (L.) Willd. *Tree Genet. Genomes* 11, 80-92.
- Doležal, J., Greilhuber, J., Suda, J. (2007). Estimation of nuclear DNA content in plants using flow cytometry. *Nature Protocols* 2, 2233-2244.
- Franks, P. J., Farquhar, G. D. (2007). The mechanical diversity of stomata and its significance in gas-exchange control. *Plant Physiology* 143, 78-87.
- Franks, P. J., Beerling, D. J. (2009). Maximum leaf conductance driven by CO<sub>2</sub> effects on stomatal size and density over geologic time. *Proceedings of the National Academy of Sciences* 106, 10343-10347.
- Hagerup, O. (1939). Studies on the significance of polyploidy III. *Deschampsia* and *Aira*. *Hereditas* 25, 185-192.
- Halverson K., Heard S. B., Nason J. D., Stireman, J.O., III. (2008). Origins, distribution, and local co-occurrence of polyploid cytotypes in *Solidago altissima* (Asteraceae). *Amer. J. Bot.* 95, 50-58.
- Hao, G. Y., Lucero, M. E., Sanderson, S. C., Zacharias, E. H., Holbrook, N. M. (2013). Polyploidy enhances the occupation of heterogeneous environments through hydraulic related trade-offs in *Atriplex canescens* (Chenopodiaceae). *New Phytol.* 197, 970-978
- Harbard, J.L., Griffin, A. R., Foster, S., Brooker, C., Kha, L. D., Koutoulis, A. (2012). Production of colchicine-induced autotetraploids as a basis for sterility breeding in *Acacia mangium* Willd.



542 Forestry 85, 427-436.

543

544 Henry, I. M., Dilkes, B. P., Young, K., Watson, B., Wu, H., Comai, L. (2005). Aneuploidy and  
 545 genetic variation in the *Arabidopsis thaliana* triploid response. *Genetics* 170, 1979-1988.

546

547 Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones and A. Jarvis, 2005. Very high resolution  
 548 interpolated climate surfaces for global land areas. *Int. J. of Climat.* 25, 1965-1978.

549

550 Husband, B. C. (2000). Constraints on polyploid evolution: a test of the minority cytotype exclusion  
 551 principle. *Proceedings of the Royal Society of London B: Biological Sciences* 2671: 217-223.

552

553 Husband, B. C. (2004). The role of triploid hybrids in the evolutionary dynamics of mixed-ploidy  
 554 populations. *Biol. J. Linn. Soc.* 82, 537-546.

555

556 Johnsson, H. (1940). Cytological studies of diploid and triploid *Populus tremula* and crosses between  
 557 them. *Hereditas* 26, 321-352.

558

559 Jørgensen, M. H., Ehrich, D., Schmickl, R., Koch, M. A., Brysting, A. K. (2011). Interspecific and  
 560 interploidal gene flow in Central European *Arabidopsis* (Brassicaceae). *BMC Evolutionary Biology*  
 561 11, 346-359.

562

563 Kennedy, B. F. Sabara H. A., Haydon, D., Husband B. C. (2006). Pollinator-mediated assortative  
 564 mating in mixed ploidy populations of *Chamerion angustifolium* (Onagraceae). *Oecologia* 150: 398-  
 565 408.

566

567 Koutecky, P., Stepanek, J., Bad'urova, T. (2012). Differentiation between diploid and tetraploid  
 568 *Centaurea phrygia*: mating barriers, morphology and geographic distribution. *Preslia* 84, 1-32.

569

570 Kudo, N., Kimura, Y. (2002). Nuclear DNA endoreduplication during petal development in cabbage:  
 571 relationships between ploidy levels and size. *J. Exp. Bot.* 53, 1017-1023.

572

573 Leitch, A. R., Leitch, I. J. (2008). Genomic plasticity and the diversity of polyploid plants. *Science*  
 574 320, 481-483.

575

576 Levin, D. A. (2002). *The role of chromosomal change in plant evolution*: Oxford University Press  
 577 New York.

578

579 Li, W. L., Berlyn, G. P., Ashton, P. M. S. (1996). Polyploids and their structural and physiological  
 580 characteristics relative to water deficit in *Betula papyrifera* (Betulaceae). *American Journal of*  
 581 *Botany* 83, 15-20.

582

583 Li, W. D., Biswas, D. K., Xu, H., Xu, C. Q., Wang, X. Z., Liu, J. K., Jiang, G. M. (2009).  
 584 Photosynthetic responses to chromosome doubling in relation to leaf anatomy in *Lonicera japonica*  
 585 subjected to water stress. *Functional Plant Biology* 36, 783-792

586

587 Liu, S., Sumei, C., Yu, C., Zhiyong, G., Dongmei, Y., Fadi, C. (2011). In vitro induced tetraploid of  
 588 *Dendranthema nankingense* (Nakai) Tzvel. shows an improved level of abiotic stress tolerance.  
 589 *Scientia Horticulturae* 127, 411-419.

590

591 Love, Á., Love, D. (1949). The geobotanical significance of polyploidy. I. Polyploidy and latitude.

Portugaliae Acta Biologica, Serie A. Morfologia, fisiologia, genetica e biologia geral. 273-352.

Lowry, E., Lester, S. E. (2006). The biogeography of plant reproduction: potential determinants of species' range sizes. *Journal of Biogeography* 33, 1975-1982.

Madlung, A. (2013). Polyploidy and its effect on evolutionary success: old questions revisited with new tools. *Heredity* 110, 99-104.

Maherali, H., Walden, A. E., Husband, B. C. (2009). Genome duplication and the evolution of physiological responses to water stress. *New Phytol.* 184, 721-731.

Manton, I. (1934). The problem of *Biscutella laevigata* L. *Molecular and General Genetics* 67, 41-57.

Manzaneda, A. J., Rey, P. J., Bastida, J. M., Weiss-Lehman, C., Raskin, E., Mitchell-Olds, T. (2012). Environmental aridity is associated with cytotype segregation and polyploidy occurrence in *Brachypodium distachyon* (Poaceae). *New Phytol.* 193, 797-805.

Martin, S. L., Husband, B. C. (2009). Influence of phylogeny and ploidy on species ranges of North American angiosperms. *Journal of Ecology* 97, 913-922.

Maslin, B. R. (2006). Generic and infrageneric names in *Acacia* following retypification of the genus. *World Wide Wattle*:1-3.

Mishra, M.K. (1997). Stomatal characteristics at different ploidy levels in *Coffea* L. *Annals Bot.* 80, 689-692.

Nobel, P.S. (2009). *Physicochemical and environmental plant physiology*. Elsevier, London. 582 pp.

O'Brien, T. P., McCully, M. E. (1981). *The study of plant structure: principles and selected methods*. Melbourne, Termarcaphi Pty. LTD.

Odee, D. W., Telford, A., Wilson, J., Gaye, A., Cavers, S. (2012). Plio-Pleistocene history and phylogeography of *Acacia senegal* in dry woodlands and savannahs of sub-Saharan tropical Africa: evidence of early colonisation and recent range expansion. *Heredity* 109, 372-382.

Odee, D.W., Wilson, J., Omondi, S., Perry, A., Cavers, S. (2015). Rangewide ploidy variation and evolution in *Acacia senegal*: a north-south divide? *AoB Plants* 7, plv011; doi:10.1093/aobpla/plv011.

Otto, F. (1990). DAPI staining of fixed cells for high-resolution flow cytometry of nuclear DNA. *Methods in Cell Biology* 33, 105-110.

Otto, S. P., Whitton, J. (2000). Polyploid incidence and evolution. *Annual Review of Genetics* 34, 401-437.

Pettigrew, F. R. S., Jack, D., Bell, K. L., Bhagwandin, A., Grinan, E., Jillani, N. (2012). Morphology, ploidy and molecular phylogenetics reveal a new diploid species from Africa in the baobab genus *Adansonia* (Malvaceae: Bombacoideae). *Taxon* 61, 1240- 1250.

641 Pontanier R., Diouf M., Zaafour M.S. (2003). Ecologie et régime hydrique de deux formations à  
642 *Acacia raddiana* au nord et au sud du Sahara (Tunisie, Sénégal). In: Grouzis, M., Le Floch, E. (Eds).  
643 Un arbre au désert: *Acacia raddiana*. Paris : IRD, 79-101.  
644

645 Prentis, P. J., Wilson, J. R. U., Dormontt, E. E., Richardson, D. M., Lowe, A. J. (2008). Adaptive  
646 evolution in invasive species. *Trends in Plant Science* 13, 288-294.  
647

648 Ramsey, J., Schemske, D. W. (1998). Pathways, mechanisms, and rates of polyploid formation in  
649 flowering plants. *Annual Review of Ecology and Systematics* 29, 467-501.  
650

651 Ross, J. H. (1979). A conspectus of the African *Acacia* species. *Memoirs of the Botanical Survey of*  
652 *South Africa* volume 44 Botanical Research Institute, Dept. of Agricultural Technical Services.  
653

654 SAS Institute Inc. 2014. SAS/STAT® 13.2 User's Guide. Cary, NC: SAS Institute Inc.  
655

656 Schlaepfer, D. R., Edwards, P. J., Billeter, R. (2010). Why only tetraploid *Solidago gigantea*  
657 (Asteraceae) became invasive: a common garden comparison of ploidy levels. *Oecologia* 163, 661-  
658 673.  
659

660 Slotte, T., Huang, H., Lascoux, M., Ceplitis, A. (2008). Polyploid speciation did not confer instant  
661 reproductive isolation in *Capsella* (Brassicaceae). *Mol. Biol. Evol* 25, 1472- 1481.  
662

663 Soltis, D. E., Soltis, P. S. (1999). The dynamic nature of polyploid genomes. *Proceedings of the*  
664 *National Academy of Sciences* 92, 8089-8091.  
665

666 Soltis, D. E., Soltis, P. S., Tate, J. A. (2004). Advances in the study of polyploidy since plant  
667 speciation. *New Phytol.* 161, 173-191.  
668

669 Soltis, D. E., Buggs, R. J. A., Doyle, J. J., Soltis, P. S. (2010). What we still don't know about  
670 polyploidy. *Taxon* 59, 1387-1403.  
671

672 Stebbins, Jr. (1950). *Variation and evolution in plants*. Columbia University Press.  
673

674 Stebbins, Jr. (1971). *Chromosomal evolution in higher plant*. Edward Arnold Ltd., London.  
675

676 Tandon, R., Shivanna, K.R. Mohan Ram, H. Y. (2001). Pollination biology and breeding system of  
677 *Acacia senegal*. *Botanical Journal of the Linnean Society* 135: 251-262.  
678

679 Van de Peer, Y., Maere, S., Meyer, A. (2009). The evolutionary significance of ancient genome  
680 duplications. *Nature Review Genetics* 10, 725-732.  
681

682 Verslues, P. E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J., Zhu, J.-K. (2006). Methods and concepts  
683 in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status.  
684 *The Plant J.* 45, 523-539.  
685

686 Wekesa, C., Makenzi, P., Chikamai, B. N., Lelon, J. K., Luvanda, A. M., Muga, M. (2009). Gum  
687 arabic yield in different varieties of *Acacia senegal* (L.) Willd in Kenya. *Afr. J. Plant Sci.* 11, 263-  
688 276.  
689

690 **Figure legends**

691  
692 **Figure 1.** Location of the progeny trial (Dahra) and the four tested provenances of *Acacia senegal* in  
693 Senegal

694  
695 **Figure 2.** Performance of polyploids expressed in percent of diploid performance. The vertical dotted  
696 line (100%) denote diploids. Error bars denote the 95% confidence limits of differences between  
697 polyploids and diploids.

698 **Figure 3.** Variation in height between diploid (filled circles) and polyploid seedlings (open circles) of  
699 *Acacia senegal* under drought stress conditions. The arrow indicates onset of the drought stress,  
700 while error bars denote SD.

701  
702  
703

Provisional

704 **Tables**

705

706 **Table 1.** F-tests for significance in growth traits between ploidy levels and provenances in the field trial and  
707 LS estimated averages of diploid and polyploid trees

Traits	Ploidy level			Provenance			Ploidy (Average performance)			
	<i>Df</i> ;	<i>F</i>	<i>P&gt;F</i>	<i>Df</i> ;	<i>F</i>	<i>P&gt;F</i>	Diploid (LSmean)	Polyploid (LSmean)	Diploid (Mean)	Polyploid (Mean)
Height 1995 (cm)	1; 29	5.96	0.015	3; 29	14.61	0.001	38.2(1.18)	45.0(2.29)	38.2 (16.3)	57.0(22.3)
Height 2012 (m)	1; 29	9.52	0.002	3; 29	4.24	0.006	4.52(0.05)	4.92(0.11)	4.57(0.05)	5.25(0.07)
Diameter 2012 (cm)	1; 29	17.29	0.001	3; 29	3.64	0.013	11.86(0.19)	13.92(0.41)	11.57(0.16)	14.34(0.28)
Crown diameter 2012 (m)	1; 29	0.06	0.804	3; 29	1.76	0.152	5.50(0.07)	5.55 (0.14)	5.40(0.06)	5.61(0.09)

708 Least square means (LS means), Simple average (Mean) and standard errors (SE) are estimated from the  
709 analysis of variance. *Df*: Degrees of freedom; *F*: *F*-value.

710

711

712

713

714 **Table 2.** F-tests for significance of morphological differences between ploidy levels and provenances of  
715 *Acacia senegal* in the growth chamber trial.

Traits	Ploidy level			Ploidy LS means	
	Df; Error	F	P>F	Diploid	Polyploid
<b>Seed traits</b>					
Number of seeds per pod	1; 55	0.34	0.56	4.4 (0.2)	4.1 (0.5)
Seed length (mm)	1; 49	13.6	0.0006	7.7 (0.1)	8.7 (0.2)
Seed width (mm)	1; 49	6.5	0.01	7.8 (0.1)	8.6 (0.2)
Germination rate (%)	1; 55	0.94	0.34	89 (3)	95 (6)
<b>Growth and morphology at 87 % field capacity</b>					
Initial total fresh weight (g)	1; 49	12.2	0.001	0.17 (0.01)	0.27 (0.02)
No. leaves	1; 48	0.02	0.89	16.0 (1.0)	16.3 (2.2)
No. branches	1; 48	0.00	0.99	6.2 (0.4)	6.2 (0.8)
Leaf length (cm)	1; 31	0.03	0.87	1.9 (0.1)	1.9 (0.2)
Leaflet length (mm)	1; 31	5.1	0.03	6.1 (0.2)	7.3 (0.5)
Leaflet width (mm)	1; 31	2.0	0.17	1.8 (0.1)	2.1 (0.1)
Stomatal density (mm <sup>-2</sup> )	1; 31	13.7	0.0008	204 (7)	139 (15)
Stomatal length (µm)	1; 31	57	<0.0001	46 (1)	71 (3)
<b>Growth at 47 % field capacity</b>					
Height at week 8 (cm)	1; 48	0.97	0.33	15.9 (0.6)	17.6 (1.4)
Height at week 9 (cm)	1; 48	2.4	0.13	17.2 (0.7)	20.0 (1.5)
Height at week 10 (cm)	1; 48	4.1	0.05	18.1 (0.7)	22.2 (1.6)
Height at week 11 (cm)	1; 48	6.3	0.02	19.1 (0.8)	24.5 (1.7)
Total fresh weight (g)	1; 48	3.6	0.06	1.0 (0.1)	1.6 (0.2)
Total dry weight (g)	1; 48	1.6	0.21	0.33 (0.03)	0.44 (0.07)
Fresh weight/dry weight ratio	1; 47	5.4	0.03	3.30 (0.05)	3.60 (0.10)

716 Least square means (LS means) and standard errors (SE) are given for the two ploidy levels. Df: Degrees of  
717 freedom and F: F-value.

718

719 **Appendix 1.** Ploidy levels in a sub-set (76 parents) of *Acacia senegal* in the progeny trial and in their offspring revealed by eight polymorphic  
720 microsatellites and flow cytometry (FCM) respectively  
721

Parent ploidy levels							Offspring ploidy levels						
Provenance	Family	Ind.	No. loci with 1-2 alleles	No. loci with more than 2 alleles	FCM on twig	No. Pods tested	No. offspring tested	Diploid (2n)	Triploid (3n)	Tetra (4n)	Penta (5n)	Hexa (6n)	Octo (8n)
Ngane	NG4	*B14	5	3	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG4	B16	4	4	Na	5	11	-	-	10	-	1	-
<b>Ngane</b>	<b>NG7</b>	<b>*B22</b>	<b>8</b>	<b>0</b>	<b>Diploid</b>	<b>1</b>	<b>2</b>	<b>2</b>	-	-	-	-	-
Ngane	NG10	*B8	7	1	Tetraploid	6	13	-	-	11	1	1	-
<b>Ngane</b>	<b>NG11</b>	<b>*B17</b>	<b>8</b>	<b>0</b>	<b>Diploid</b>	<b>1</b>	<b>2</b>	<b>2</b>	-	-	-	-	-
Ngane	NG14	B1	5	3	Na	5	12	-	-	11	-	1	-
Ngane	NG14	*B2	5	3	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG15	*B16	5	3	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG16	B3	5	3	Na	4	9	-	-	9	-	-	-
Ngane	NG16	*B19	4	4	Tetraploid	1	2	-	-	1	-	1	-
Ngane	NG17	*B8	5	3	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG18	*B25	4	4	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG19	B1	4	4	Na	4	8	-	-	6	-	2	-
Ngane	NG19	* B24	4	4	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG20	*B3	4	4	Tetraploid	6	16	-	-	15	-	-	1
Ngane	NG21	* B4	4	4	Tetraploid	6	13	-	-	12	-	1	-
Ngane	NG22	*B6	5	3	Tetraploid	1	2	-	-	2	-	-	-
Ngane	NG25	*B30	4	0	Diploid	4	10	10	-	-	-	-	-
Ngane	NG26	*B17	4	4	Tetraploid	1	2	-	-	2	-	-	-
Diamenar	DIA2	B1	8	0	Na	1	6	6	-	-	-	-	-
Diamenar	DIA2	*B2	8	0	Diploid	1	2	2	-	-	-	-	-
Diamenar	DIA6	*B2	8	0	Diploid	1	2	2	-	-	-	-	-
Diamenar	DIA6	B3	8	0	Na	1	4	4	-	-	-	-	-
Diamenar	DIA7	*B27	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA8	*B21	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA11	* B5	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA13	*B26	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA14	*B6	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA15	B8	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA15	*B21	8	0	Diploid	1	2	2	-	-	-	-	-
Diamenar	DIA17	*B17	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA18	B4	8	0	Diploid	1	2	2	-	-	-	-	-
Diamenar	DIA20	*B23	8	0	Na	1	2	2	-	-	-	-	-

Diamenar	DIA22	*B6	8	0	Na	1	3	3	-	-	-	-	-
Diamenar	DIA22	*B14	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA26	*B2	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA27	*B5	8	0	Na	1	2	2	-	-	-	-	-
Diamenar	DIA29	*B5	8	0	Na	1	2	2	-	-	-	-	-
Daiba	DA1	B17	7	1	Triploid	1	3	-	2	1	-	-	-
Daiba	DA1	*B19	8	0	Diploid	1	5	5	-	-	-	-	-
Daiba	DA2	*B5	8	0	Diploid	1	3	3	-	-	-	-	-
Daiba	DA4	*B12	8	0	Diploid	1	3	3	-	-	-	-	-
Daiba	DA4	*B22	7	1	Na	1	3	-	-	3	-	-	-
Daiba	DA6	*B6	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA7	*B1	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA8	*B17	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA13	*B9	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA15	*B3	7	1	Tetraploid	1	2	-	-	-	-	2	-
Daiba	DA15	B11	8	0	Diploid	1	5	5	-	-	-	-	-
Daiba	DA16	*B15	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA17	*B2	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA18	B2	8	0	Diploid	1	4	4	-	-	-	-	-
Daiba	DA18	B5	8	0	Diploid	1	1	1	-	-	-	-	-
Daiba	DA19	*B6	8	0	Diploid	1	2	2	-	-	-	-	-
Daiba	DA20	*B13	8	0	Diploid	1	4	4	-	-	-	-	-
Daiba	DA25	*B8	8	0	Diploid	1	3	3	-	-	-	-	-
Daiba	DA25	B12	8	0	Diploid	1	3	3	-	-	-	-	-
Daiba	DA26	B2	8	0	Diploid	1	5	5	-	-	-	-	-
Daiba	DA26	B9	8	0	Diploid	1	3	3	-	-	-	-	-
Kidira	K1	*B3	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K3	*B2	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K4	*B23	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K5	*B7	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K7	*B8	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K8	*B7	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K9	*B5	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K14	*B6	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K16	*B15	8	0	Diploid	1	4	4	-	-	-	-	-
Kidira	K17	B14	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K20	*B9	8	0	Diploid	1	4	4	-	-	-	-	-
Kidira	K21	B13	8	0	Diploid	1	5	5	-	-	-	-	-
Kidira	K21	*B22	4	4	Tetraploid	1	2	-	-	2	-	-	-
Kidira	K22	*B9	8	0	Diploid	1	2	2	-	-	-	-	-
Kidira	K23	B3	6	2	Na	1	Na	Na	-	Na	-	-	-



	Kidira	K23	B6	8	0	Diploid	1	4	4	-	-	-	-	-
	Kidira	K25	*B16	4	4	Tetraploid	1	2	-	-	2	-	-	-
722	In bold: The pure diploid Ngane families (NG7 & NG11). Dark grey: triploid mother producing both triploid and tetraploid offspring. Light grey: Families with mixed													
723	ploidy levels. Individuals marked with * were used in the drought stress trial. Na: Not assessed.													
724														

Provisional

Figure 01.JPEG




Figure 02.TIF




Figure 03.JPEG

